

Dissemination of an AbaR-type Resistance Island in Multidrug-resistant *Acinetobacter baumannii* Global Clone 2 in Daejeon of Korea

Ji Youn Sung¹, Sun Hoe Koo², Hye Hyun Cho³, Kye Chul Kwon²

¹Department of Biomedical Laboratory Science, Far East University, Eumseong,

²Department of Laboratory Medicine, College of Medicine, Chungnam National University, Daejeon,

³Department of Biomedical Laboratory Science, Jeonju Kijeon College, Jeonju, Korea

Background: *Acinetobacter baumannii* resistance islands (AbaRs) are transposons that have the role of important vehicles for the acquisition of antimicrobial resistance genes, and are associated with multidrug resistance (MDR). In this study, we aimed to determine the AbaRs in MDR *A. baumannii* global clone 2 (GC2) clinical isolates obtained from a university hospital in Daejeon, Korea.

Methods: This study included 17 MDR *A. baumannii* strains isolated in Daejeon, Korea. The minimal inhibitory concentrations (MICs) were determined by Etest. *A. baumannii* isolates were characterized using 2 multiplex PCR assays and a multilocus sequence typing (MLST) scheme. To detect and characterize AbaRs, PCR and PCR mapping experiments were performed.

Results: All 17 MDR *A. baumannii* isolates tested in

this study belonged to GC2 and contained 5 sequence types (STs): 75, 92, 137, 138, and 357. Tn6166 that contains antimicrobial resistance genes and is also known as AbaR4a was found in all 17 GC2 strains. This is the first report of Tn6166 in MDR *A. baumannii* GC2 isolates in Korea. In contrast, AbaR4 was not found in the GC2 isolates.

Conclusion: Tn6166 has been disseminated among MDR *A. baumannii* GC2 isolates in Korea. Further investigation is needed to recover the various types of AbaRs in MDR *A. baumannii* GC2 isolates in Korea are responsible for the multiple antimicrobial resistance mechanisms. (**Ann Clin Microbiol 2013;16: 75-80**)

Key Words: *A. baumannii*, Transposon, Multilocus sequence typing, PCR

INTRODUCTION

Multidrug-resistant (MDR) *A. baumannii* strains have been reported worldwide usually belonged to one of two globally distributed clonal complexes. These two clones are global clones 1 and 2 (GC1 and GC2) which are also known as European clone I and II, respectively [1]. Most MDR *A. baumannii* isolates belonging to GC1 and GC2 are important opportunistic pathogen associated with nosocomial infections and hospital outbreaks. The evolution of MDR *A. baumannii* strains have been suggested to decrease cell wall permeability, increase constitutive expression of efflux pumps and/or acquire their antimicrobial resistance genes via integrons, transposons and plasmid [2]. In particular, rapid dissemination of acquired resistance

genes has led to the widespread of MDR *A. baumannii* strains which give rise to significant therapeutic failures and increase of patient morbidity and mortality [3].

Many MDR *A. baumannii* strains harbor multiple antimicrobial resistance genes, which are integrated in the chromosomal *comM* gene encoding an ATPase domain. Such genomic resistance islands were designated *A. baumannii* resistance islands (AbaRs) [3]. AbaRs are transposons that incorporate themselves into the *comM* gene and have role as important vehicles for the acquisition of antimicrobial resistance genes. The first example of AbaRs was found in strain AYE and designated AbaR1 island, which includes more than 40 antimicrobial and heavy metal resistance genes. Until recently, many types of AbaRs have been fully characterized following detection of AbaR1 and most of them belong to AbaR3-type and AbaR4-type islands [3,4].

AbaR3-type islands found in GC1 isolates consist of a backbone transposon, Tn6019, which interrupted by a large compound transposon containing a variable resistance region located

Received 12 September, 2012, Revised 21 November, 2012

Accepted 10 December, 2012

Correspondence: Sun Hoe Koo, Department of Laboratory Medicine, College of Medicine, Chungnam National University, 640, Daesadong, Jung-gu, Daejeon 301-721, Korea. (Tel) 82-42-280-7798, (Fax) 82-42-257-5365, (E-mail) azaza72@naver.com

between two copies of transposon, Tn6018 [4]. AbaR4-type islands found in both GC1 and GC2 isolates are consisted of a backbone transposon, Tn6022, which is most closely related to the transposon Tn6021 that interrupts the *comM* gene but carries no resistance genes [5,6]. Tn6022 differs from Tn6021 in that it includes a short different patch of 90.8% identity in the *tniC-tniA* gene region (shown as a black line in Fig. 1) and lacks of *tniD* gene and part of *tniB* and *tniE* genes [4].

Even though various AbaRs have been recovered in MDR *A. baumannii* isolates worldwide [6,7], there is a relative paucity of reports about AbaRs in MDR *A. baumannii* isolates from Korea. In the present study, we aimed to investigate the AbaRs for understanding of acquired antimicrobial resistances mechanisms in MDR *A. baumannii* GC2 isolates obtained from a university hospital in Daejeon, Korea.

MATERIALS AND METHODS

1. Selection of bacteria isolates

Seventeen MDR *A. baumannii* isolates were collected and characterized in our previous study [8]. The isolates were col-

lected from different patients in one university hospital in Daejeon (Korea) during the period between 2009 and 2011. MDR phenotypes were defined as showing resistance to representative antimicrobial agents of at least three different classes of drugs: aminoglycosides (gentamicin, amikacin), antipseudomonal penicillins (ticarcillin, piperacillin, piperacillin/tazobactam), carbapenems (imipenem, meropenem), antipseudomonal cephalosporins (ceftazidime, cefepime), and fluoroquinolone (ciprofloxacin) [9].

2. DNA extraction and PCR amplification

Whole-cell (genomic) DNA was obtained from each target strain by using a genomic DNA purification kit (SolGent, Daejeon, Korea) according to the manufacturer's instructions. PCR was performed using 50 ng of genomic DNA, 2.5 μ L of 10 \times Taq buffer, 0.5 μ L of 10 mM dNTP mix, 20 pmol of each primer, and 0.7 U of Taq DNA polymerase (SolGent), in a total volume of 25 μ L. Each target site was amplified in a GeneAmp PCR System 9600 thermal cycler (Perkin-Elmer Cetus Corp., Norwalk, CT, USA). The amplified products were separated via electrophoresis on 1.5% (w/v) agarose gels containing ethidium




Fig. 1. Schematic representation of transposons isolated from multidrug-resistant *A. baumannii* strains. The dotted rectangle in Tn6019 represents deletion part in Tn6021. A short patch of 90.8% identity in the *tniC-tniA* gene region was shown as a black line in Tn6022 and Tn6166. In Tn6022, the dotted rectangle exhibits deletion part in Tn6166. The horizontal arrows indicate the translation orientation of the genes.

bromide, and visualized using a BioDoc-14TM Imaging system (UVP, Cambridge, UK). For sequencing, PCR products were purified with a PCR purification kit (SolGent) following the manufacturer’s protocols.

3. Characterization of *A. baumannii* isolates

The two multiplex PCR assays were used as previously described [10] to identify members of the GC1 and GC2. The Oxford multilocus sequence typing (MLST) scheme [11], which

uses seven housekeeping genes (*gltA*, *gyrB*, *gdhB*, *recA*, *cpn60*, *gpi* and *rpoD24*) was used to determine the sequence types (STs). A ST number was assigned by comparing the allele sequences to ones in the MLST site (<http://pubmlst.org/abaumannii/>).

4. Detections and characterizations of AbaRs

The genes associated with AbaRs were detected by PCR using the published primers (Table 1) [2,3]. The amplification regions are represented in Fig. 2. To investigate the structures of

Table 1. Primer pairs used for the detection and characterization of *A. baumannii* resistance islands

Region	Primer	Sequence 5’-3’	Amplicon length (bp)
<i>comM</i>	RH791 RH913	TGCTGCAATGAGCTGAAAAGT GCCTCTCATTGAGGTTGAGG	982
<i>comM</i> -AbaR (J1)	RH927 RH792	CAACCCTGTCTTTGCATTTG TTCGAGCTTGAAAACGTGCAC	846
AbaR- <i>comM</i> (J2)	RH916 RH928	CCCAAATACTGCCATGTTGA GCCAGCAAGCTCAGCATAA	796
<i>Tn6018</i>	RH768 RH769	GAATCGCTGGTGATGATGGC GGTCTGAGACTTCGTGAGCGC	1,630
<i>uspA</i>	RH919 RH793	TGTCAAAAATTATTGCATGT CCCAAGAGAGCTGATTTTGC	632
<i>comM</i> - <i>tniB</i> Δ	RH791 RH909	TGCTGCAATGAGCTGAAAAGT GCGATTCAAAAATATCGGTCAA	3,119
<i>comM</i> -orf/ <i>uspA</i>	RH791 RH592	TGCTGCAATGAGCTGAAAAGT AAGCTTATCGAAAAGGCGTAGA	4,632
<i>tniB</i> - <i>tniE</i>	RH910 RH587	GCGATAGTGAACGGATTGAGA TTGCCATTAAGCACAACAG	560
<i>tniB</i> Δ- <i>sup</i>	RH910 RH771	GCGATAGTGAACGGATTGAGA TGTA AAAATCTGGTGGTCGTAC	3,267
<i>sup</i> - <i>tetA</i> (B)	RH772 tetBF	GCAGCCATAGGAATGACTTTTA TTGGTTAGGGGCAAGTTTTG	3,961
orf4-CR2	RH916 LECR2	CCCAAATACTGCCATGTTGA CACTGGCTGGCAATGTCTAG	4,466
CR2- <i>comM</i>	RH702 RH928	CTCGTCAACGATCTGATAGAGAAGGG GCCAGCAAGCTCAGCATAA	5,021




Fig. 2. Schematic representation of transposon, Tn6166 and the amplification regions used in the diagnostic PCRs. The genes are shown by horizontal arrows with the gene name above. The dotted lines indicate PCR amplification regions with primer name on the right. Abbreviation: Tn, transposon.

Table 2. Properties of multidrug-resistant *A. baumannii* GC2 isolates carrying Tn6166

Sequence type	Number of isolates	Global clone	<i>bla</i> _{OXA-51} like	Island junctions				Features of Tn6166				
				<i>comM</i>	<i>comM</i> -AbaR J1	AbaR- <i>comM</i> J2	Tn6018	<i>uspA</i>	<i>sul</i>	<i>strA</i> , <i>strB</i>	<i>tetB</i> , <i>tetR</i>	ISCR2
75	1	2	OXA-66	-	+	+	-	+	+	+	+	+
92	2	2	OXA-66	-	+	+	-	+	+	+	+	+
137	7	2	OXA-66	-	+	+	-	+	+	+	+	+
138	3	2	OXA-66	-	+	+	-	+	+	+	+	+
357	4	2	OXA-66	-	+	+	-	+	+	+	+	+

Abbreviation: AbaR, *A. baumannii* resistance island.

AbaR backbone, PCR mapping experiments were performed as described previously [2,3]. The amplicons were purified and sequenced using a BigDye Terminator Cycle Sequencing Kit (PE Applied Biosystems, Foster City, CA, USA) and an ABI PRISM 3730XL DNA analyzer (PE Applied Biosystems). DNA fragments (up to 1 kb in size) were sequenced using the overlapping PCR technique. The various DNA sequences were confirmed by using the BLAST paired alignment facility (<http://blast.ncbi.nlm.gov>).

RESULTS

1. Characterization of MDR *A. baumannii*

Association of the GC lineages was tested for 17 MDR *A. baumannii* using multiplex PCRs. All 17 isolates tested in this study belonged to GC2 and carried allele 66 of the intrinsic *bla*_{OXA-51}-like genes which corresponds with their assignment to GC2. MLST analysis of GC2 isolates revealed 5 STs (75, 92, 137, 138 and 357) (Table 2).

2. MDR *A. baumannii* GC2 strains including AbaRs

To determine whether MDR *A. baumannii* GC2 strains harbored a transposon in *comM* gene, all 17 strains were tested for the presence of *comM* gene. An amplicon of *comM* gene was not obtained from any of our isolates except *Acinetobacter calcoaceticus* control strain. J1 (*comM*-AbaR) and J2 (AbaR-*comM*) segments, which form the boundaries of the AbaR backbone transposon, Tn6019, were amplified in all 17 isolates, indicating that transposon related insertions are present at the position of *comM* gene. However, Tn6018 and an interrupted *uspA* gene were not present in all isolates tested in present study. These results indicated that AbaR3-type islands were not present in GC2 isolates.

3. Characterization of AbaRs

To characterize AbaRs contained in GC2 isolates, we mapped the continuous regions from J1 to J2 segment by overlapping PCRs and the sequencing. The about 18 kb transposon, Tn6166 (GenBank accession no. JN247441), was identified from the genome of all MDR *A. baumannii* isolates which were tested in this study and belonged to GC2 (Fig. 1). Tn6166 carries Tn6022 containing intact universal stress protein (*uspA*) and sulfate permease (*sul*) except for a 2.85 kb deletion region which removes *tniD* gene and part of *tniB* and *tniE* genes. The *tet(B)* determinant and *strA* and *strB* genes are located in Tn6166 harboring the small mobile element CR2.

DISCUSSION

MDR *A. baumannii* isolates pose therapeutic problems caused by intrinsic antimicrobial resistance mechanisms and capability to acquire multiple antimicrobial resistance genes via integron and transposon. Many researchers tried to find the role of integron and transposon mediated resistance in developing MDR *A. baumannii* strains, and recently recognized AbaRs as transposons containing multiple resistance determinants [2]. Further studies have found the frequent occurrence and worldwide dissemination of various AbaRs among MDR *A. baumannii* isolates [12]. However, much less is known about AbaRs in MDR *A. baumannii* isolates in Korea.

In this study, structure of transposon in MDR *A. baumannii* GC2 isolates was determined by using a strategy based on the structural and sequence homology of the AbaRs [2,3]. All GC2 isolates harbored an interrupted *comM* gene and uninterrupted *uspA* gene. In addition, Tn6018 was not found in any isolates tested present study. Our results suggest that transposon insertion in GC2 isolates was not closely connected to the

Tn6018-containing AbaR3-type islands found in most of GC1 isolates. To characterize the transposons we used published primers for Tn6166, which is a type of genomic resistance island found in GC2 isolates and also known as AbaR4a for the structural similarity to Tn6022, the backbone of AbaR4 [13]. The Tn6166 was found in all 17 MDR *A. baumannii* GC2 strains which were collected during the three years and revealed various ST types (ST 75, 92, 137, 138 and 357). This result suggests that Tn6166 was disseminated in GC2 isolates and have existed for many years, though there have not been any reports as yet concerning Tn6166 in Korea. This is the first report of Tn6166 in MDR *A. baumannii* GC2 isolates in Korea.

Tn6166 was first detected in *A. baumannii* RUH134/A320 which was isolated in 1982 in the Netherlands and has been used as the reference strain for GC2 isolates [4]. This suggests that Tn6166 has been served as a reservoir for antimicrobial resistance genes in GC2 isolates since long before. The genomic island has a different origin to the AbaR3-type islands found in GC1 isolates and representative of a type of genomic island found in GC2 isolates.

Reportedly, AbaR4 which carries transposon Tn2006 harboring the *bla*_{OXA-23} carbapenem resistance gene have been identified in GC1 and GC2 isolates [14,15]. AbaR4 globally disseminated has been recovered in Australia, UK and Taiwan [5, 16,17], but there is not yet any data about AbaR4 in Korea so far. Similarly, in our examination, AbaR4 was not found in any isolates, although the 5 isolates of them harbored *bla*_{OXA-23} gene.

In present study, it was found that various MDR *A. baumannii* GC2 isolates collected during the period between 2009 and 2011 contained Tn6166 containing antimicrobial resistance genes. This result suggests that Tn6166, AbaR4-type island-related transposons, has been disseminated among MDR *A. baumannii* GC2 isolates in Korea before the transposons were determined in GC2 isolates. This finding emphasizes that further investigation is needed to recover various AbaRs in MDR *A. baumannii* GC2 isolates in Korea because AbaRs may seem to play a substantial role in antimicrobial resistance in MDR *A. baumannii* isolates through insertion and homologous recombination of various antimicrobial resistance genes.

REFERENCES

1. Hamouda A, Evans BA, Towner KJ, Amyes SG. Characterization of epidemiologically unrelated *Acinetobacter baumannii* isolates from four continents by use of multilocus sequence typing, pulsed-field gel electrophoresis, and sequence-based typing of *bla*_{OXA-51}-like genes. *J Clin Microbiol* 2010;48:2476-83.
2. Petersen A, Guardabassi L, Dalsgaard A, Olsen JE. Class I integrons containing a *dhfr*I trimethoprim resistance gene cassette in aquatic *Acinetobacter* spp. *FEMS Microbiol Lett* 2000;182:73-6.
3. Post V, White PA, Hall RM. Evolution of AbaR-type genomic resistance islands in multiply antibiotic-resistant *Acinetobacter baumannii*. *J Antimicrob Chemother* 2010;65:1162-70.
4. Nigro SJ and Hall RM. Antibiotic resistance islands in A320 (RUH134), the reference strain for *Acinetobacter baumannii* global clone 2. *J Antimicrob Chemother* 2012;67:335-8.
5. Hamidian M and Hall RM. AbaR4 replaces AbaR3 in a carbapenem-resistant *Acinetobacter baumannii* isolate belonging to global clone 1 from an Australian hospital. *J Antimicrob Chemother* 2011;66:2484-91.
6. Kim DH, Choi JY, Jung SI, Thamlikitkul V, Song JH, Ko KS. AbaR4-type resistance island including the *bla*_{OXA-23} gene in *Acinetobacter nosocomialis* isolates. *Antimicrob Agents Chemother* 2012;56:4548-9.
7. Lee HY, Chang RC, Su LH, Liu SY, Wu SR, Chuang CH, et al. Wide spread of Tn2006 in an AbaR4-type resistance island among carbapenem-resistant *Acinetobacter baumannii* clinical isolates in Taiwan. *Int J Antimicrob Agents* 2012;40:163-7.
8. Sung JY, Kwon KC, Cho HH, Koo SH. Antimicrobial resistance determinants in imipenem-nonsusceptible *Acinetobacter calcoaceticus-baumannii* complex isolated in Daejeon, Korea. *Korean J Lab Med* 2011;31:265-70.
9. Lee K, Yong D, Jeong SH, Chong Y. Multidrug-resistant *Acinetobacter* spp.: increasingly problematic nosocomial pathogens. *Yonsei Med J* 2011;52:879-91.
10. Turton JF, Gabriel SN, Valderrey C, Kaufmann ME, Pitt TL. Use of sequence-based typing and multiplex PCR to identify clonal lineages of outbreak strains of *Acinetobacter baumannii*. *Clin Microbiol Infect* 2007;13:807-15.
11. Bartual SG, Seifert H, Hippler C, Luzon MA, Wisplinghoff H, Rodríguez-Valera F. Development of a multilocus sequence typing scheme for characterization of clinical isolates of *Acinetobacter baumannii*. *J Clin Microbiol* 2005;43:4382-90.
12. Bonnin RA, Poirel L, Nordmann P. AbaR-type transposon structures in *Acinetobacter baumannii*. *J Antimicrob Chemother* 2012; 67:234-6.
13. Seputiene V, Povilonis J, Suziedeliene E. Novel variants of AbaR resistance islands with a common backbone in *Acinetobacter baumannii* isolates of European clone II. *Antimicrob Agents Chemother* 2012;56:1969-73.
14. Adams MD, Goglin K, Molyneaux N, Hujer KM, Lavender H, Jamison JJ, et al. Comparative genome sequence analysis of multidrug-resistant *Acinetobacter baumannii*. *J Bacteriol* 2008;190: 8053-64.
15. Shaikh F, Spence RP, Levi K, Ou HY, Deng Z, Towner KJ, et al. ATPase genes of diverse multidrug-resistant *Acinetobacter baumannii* isolates frequently harbour integrated DNA. *J Antimicrob Chemother* 2009;63:260-4.
16. Turton JF, Baddal B, Perry C. Use of the accessory genome for characterization and typing of *Acinetobacter baumannii*. *J Clin Microbiol* 2011;49:1260-6.
17. Lee HY, Chang RC, Su LH, Liu SY, Wu SR, Chuang CH, et al. Wide spread of Tn2006 in an AbaR4-type resistance island among carbapenem-resistant *Acinetobacter baumannii* clinical isolates in Taiwan. *Int J Antimicrob Agents* 2012;40:163-7.

=국문초록=

대전에서 분리된 다제내성 *Acinetobacter baumannii* Global Clone 2에서 AbaR-type Resistance Island의 확산

¹극동대학교 임상병리학과, ²충남대학교 의과대학 진단검사의학교실, ³전주기전대학 임상병리과
성지연¹, 구선희², 조혜현³, 권계철²

배경: *Acinetobacter baumannii* resistance islands (AbaRs)는 *A. baumannii*가 항균제 내성 유전자를 획득하고 다제내성을 갖는데 중요한 역할을 하는 transposons이다. 따라서 본 연구에서는 대전의 한 대학병원에서 분리된 다제내성 *A. baumannii* global clone (GC) 2에서 AbaRs를 검출하여 항균제 내성기전을 분석하고자 하였다.

방법: 대전의 한 대학병원에서 분리된 다제내성 *A. baumannii* 17균주를 대상으로 항균제에 대한 최소발육저지농도를 결정하기 위해 Etest를 수행하였다. 또한 다중중합효소연쇄반응 및 multilocus sequence typing (MLST)을 이용하여 *A. baumannii*의 분자유전학적 특징을 결정하였다. AbaRs는 PCR mapping과 염기서열 분석을 통하여 확인하였다.

결과: 본 연구에서 분리된 다제내성 *A. baumannii* 17균주는 모두 GC2에 속하는 것으로 나타났다. 또한 이들 균주는 모두 AbaR4a로도 알려져 있는 Tn6166을 포함하고 있음이 확인되었다. 국내에서는 아직까지 Tn6166을 포함하고 있는 *A. baumannii*에 대한 보고가 없었으며 본 연구가 최초의 보고이다.

결론: 본 연구에서 Tn6166이 한국에서 분리된 다제내성 *A. baumannii* 사이에 확산되어 있음이 확인되었다. 따라서 다제내성 *A. baumannii*의 내성기전을 밝히기 위해서는 다양한 종류의 AbaRs에 대한 추가적인 연구가 필요할 것으로 사료된다.

[Ann Clin Microbiol 2013;16:75-80]

교신저자 : 구선희, 301-721, 대전시 중구 대사동 640
충남대학교병원 진단검사의학과
Tel: 042-280-7798, Fax: 042-257-5365
E-mail: azaza72@naver.com